



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

In any ellipse the product of the perpendiculars from the foci on any tangent is constant, *i. e.*, $EF \times E'F' = GF \times G'F' = \text{constant}$.

Hence $E'F'/G'F' = GF/EF = \text{constant}$, and the locus of F' is a fixed line SF' .

Now the lines PS, PF, PN, PF' form an harmonic pencil, and L, F, N, F' are harmonic points.

Hence the lines SL, SF, SN, SF' form an harmonic pencil.

But SL, SF, SF' are fixed. Therefore, SN is a fixed line.


Likewise the foot of the normal corresponding to the other tangent SP' lies on a fixed line which is the harmonic conjugate of SP' with respect to SF and SF' .

431. Proposed by E. M. MORGAN, Dartmouth College.

Trisect the angles of the triangle ABC and let the trisectors nearest each side meet in the respective points M, N, P . Prove by trigonometry that the triangle MNP is equilateral.

SOLUTION BY A. M. HARDING, University of Arkansas.

Let a, b, c denote the sides BC, CA, AB respectively. Construct the points M_1 and M_2 on CA and AB respectively, such that $CM_1 = CM$ and $BM_2 = BM$. Then $MN = M_1N$ and $MP = M_2P$.



If $3\alpha, 3\beta, 3\gamma$ represent the angles A, B, C we have

$$\overline{M_1N}^2 = \overline{AN}^2 + \overline{AM_1}^2 - 2AN \times AM_1 \cos \alpha,$$

or

$$\overline{MN}^2 = \overline{AN}^2 + (b - CM)^2 - 2AN(b - CM) \cos \alpha.$$

Also

$$\overline{PN}^2 = \overline{AN}^2 + \overline{AP}^2 - 2AN \times AP \cos \alpha.$$

Hence

$$\begin{aligned} \overline{MN}^2 - \overline{PN}^2 &= (b - CM)^2 - \overline{AP}^2 - 2AN(b - CM) \cos \alpha + 2AN \times AP \cos \alpha, \\ &= (b - CM - AP)(b - CM + AP - 2AN \cos \alpha). \end{aligned}$$

Now

$$CM = \frac{\sin \beta}{\sin (\beta + \gamma)} a, \quad AP = \frac{\sin \beta}{\sin (\alpha + \beta)} c, \quad AN = \frac{\sin \gamma}{\sin (\alpha + \gamma)} b.$$

Also

$$b = \frac{\sin 3\beta}{\sin 3\alpha} a, \quad c = \frac{\sin 3\gamma}{\sin 3\alpha} a.$$

Hence

$$\begin{aligned}
 & b - 2AN \cos \alpha - CM + AP \\
 &= \left[1 - \frac{2 \sin \gamma \cos \alpha}{\sin (\alpha + \gamma)} \right] b + \frac{\sin \beta}{\sin (\alpha + \beta)} c - \frac{\sin \beta}{\sin (\beta + \gamma)} a \\
 &= \frac{\sin (\alpha - \gamma)}{\sin (\alpha + \gamma)} b + \frac{\sin \beta}{\sin (\alpha + \beta)} c - \frac{\sin \beta}{\sin (\beta + \gamma)} a \\
 &= \left[\frac{\sin (\alpha - \gamma) \sin 3\beta}{\sin (\alpha + \gamma) \sin 3\alpha} + \frac{\sin \beta \sin 3\gamma}{\sin (\alpha + \beta) \sin 3\alpha} - \frac{\sin \beta}{\sin (\beta + \gamma)} \right] a \\
 &= \left[\frac{\sin (\alpha - \gamma) \sin 3\beta}{\sin (\alpha + \gamma)} + \frac{\sin \beta \sin 3\gamma}{\sin (\alpha + \beta)} - \frac{\sin \beta \sin 3\alpha}{\sin (\beta + \gamma)} \right] \frac{a}{\sin 3\alpha} \\
 &= \{ \sin (\alpha - \gamma)[3 - 4 \sin^2 (\alpha + \gamma)] + \sin \beta[3 - 4 \sin^2 (\alpha + \beta)] \\
 &\quad - \sin \beta[3 - 4 \sin^2 (\beta + \gamma)] \} a / \sin 3\alpha \\
 &= \{ \sin (\alpha - \gamma)[3 - 4 \sin^2 (\alpha + \gamma)] - 4 \sin \beta[\sin^2 (\alpha + \beta) \\
 &\quad - \sin^2 (\beta + \gamma)] \} a / \sin 3\alpha \\
 &= \{ \sin (\alpha - \gamma)[3 - 4 \sin^2 (\alpha + \gamma)] - 2 \sin \beta[\cos 2(\beta + \gamma) \\
 &\quad - \cos 2(\alpha + \beta)] \} a / \sin 3\alpha \\
 &= \{ \sin (\alpha - \gamma)[3 - 4 \sin^2 (\alpha + \gamma)] - 4 \sin \beta[\sin (\alpha - \gamma) \sin (\alpha + 2\beta + \gamma)] \} \\
 &\quad a / \sin 3\alpha \\
 &= \sin (\alpha - \gamma)[3 - 4 \sin^2 (\alpha + \gamma) - 4 \sin \beta \sin (\alpha + 2\beta + \gamma)] a / \sin 3\alpha \\
 &= \sin (\alpha - \gamma)[1 + 2 \cos 2(\alpha + \gamma) + 2 \cos (\alpha + 3\beta + \gamma) \\
 &\quad - 2 \cos (\alpha + \beta + \gamma)] a / \sin 3\alpha.
 \end{aligned}$$

Now

$$\cos (\alpha + 3\beta + \gamma) = -\cos 2(\alpha + \gamma)$$

and

$$\cos (\alpha + \beta + \gamma) = \cos 60^\circ = \frac{1}{2}.$$

Hence $b - 2AN \cos \alpha - CM + AP = 0$, and $MN = PN$.

Likewise $MP = PN$.

Also solved by T. M. BLAKESLEE.

CALCULUS.

335. Proposed by W. R. LEBOLD, Cambridge, Ohio.

Let $\rho = F(\theta, \phi)$ be the equation in polar coordinates of a closed surface. Show that the volume of the solid bounded by the surface is equal to the double integral

$$\frac{1}{3} \iint \rho \cos \gamma d\sigma$$